

MODULI SPACES OF 6 AND 7-DIMENSIONAL COMPLETE INTERSECTIONS

JIANBO WANG

ABSTRACT. This paper proves the existence of homeomorphic (diffeomorphic) complex 6-dimensional (7-dim) complete intersections that belong to components of the moduli space of different dimensions. These results are given as a supplement to earlier result on 5-dimensional complete intersections.

1. INTRODUCTION

Let $X_n(\underline{d}) \subset \mathbb{C}P^{n+r}$ be a smooth complete intersection of multidegree $\underline{d} = (d_1, \dots, d_r)$. The product $d_1 d_2 \cdots d_r$ is called the total degree, denoted by d .

Libgober and Wood ([8]) show the existence of homeomorphic complete intersections of dimension 2 and diffeomorphic ones of dimension 3 which belong to components of the moduli space having different dimensions. P. Brückmann ([1, 2]) shows that there are families of arbitrary length k of complete intersections in $\mathbb{C}P^{4k-2}$ and $\mathbb{C}P^{6k-2}$ (resp. $\mathbb{C}P^{5k-2}$) consisting of homeomorphic complete intersections of dimension 2 and 4 (resp. diffeomorphic ones of dimension 3) but that belong to components of the moduli space of different dimensions. By the help of Theorem 1.1 in [4], the author generalized the complex dimension of complete intersections to dimension 5 ([10]).

The goal of this paper is to give the following theorem, which is a supplement to the results of Brückmann [1, 2] and the author [10].

Theorem 1.1. *For each integer $k > 1$, there exist k homeomorphic (diffeomorphic) complex 6-dimensional (7-dim) complete intersections in $\mathbb{C}P^{8k-2}$ ($\mathbb{C}P^{15k-8}$) belonging in the moduli space to components with different dimensions.*

Acknowledgement. The author would like to thank Guo Xianqiang for his warm email conversation.

2. MODULI SPACES OF COMPLEX 6 AND 7 DIMENSIONAL COMPLETE INTERSECTIONS

Let $X_n(\underline{d}) \subset \mathbb{C}P^N$, where $n \geq 2$, $\underline{d} = (d_1, \dots, d_r)$, $d_i \geq 2$ and $r = N - n$. As usual, define the power sums $s_i = \sum_{j=1}^r d_j^i$ for $1 \leq i \leq n$, then the Pontrjagin numbers and Euler characteristic of $X_n(\underline{d})$ depend only on the dimension n , total degree d and power sums s_i ([7, §7]). Assume that $X_n(\underline{d})$ is not a $\mathbf{K}3$ -surface or a quadratic hypersurface, then from

The author is supported by NSFC grant No.11001195 and Beiyang Elite Scholar Program of Tianjin University, No.60301016.

[1, Lemma 3], the explicit formula for moduli space dimension is

$$(2.1) \quad m(\underline{d}) := m(X_n(\underline{d})) = 1 - (N+1)^2 + \sum_{i=1}^r \binom{N+d_i}{N} \\ + \sum_{i=1}^r \sum_{j=1}^r (-1)^j \sum_{1 \leq k_1 < \dots < k_j \leq r} \binom{N+d_i-d_{k_1}-\dots-d_{k_j}}{N}.$$

Where $\binom{m}{N} = 0$ for $m < N$ ($m \in \mathbb{Z}$).

Theorem 2.1. *For each integer $k > 1$, there exist k homeomorphic complex 6-dimensional complete intersections in $\mathbb{C}P^{8k-2}$ belonging in the moduli space to components with different dimensions.*

Proof. Consider the following two multidegrees([3]¹)

$$(2.2) \quad \underline{d} = (2323, 2241, 2231, 2117, 2079, 1957, 1953, 1899), \\ \underline{d}' = (2321, 2263, 2187, 2163, 2037, 2001, 1919, 1909).$$

They have the same total degree d , power sums $s_1, \dots, s_6$ as follows:

$$\begin{aligned} d &= 371\,008\,634\,983\,489\,635\,445\,991\,601, \\ s_1 &= 16\,800, \\ s_2 &= 35\,449\,960, \\ s_3 &= 75\,160\,663\,200, \\ s_4 &= 160\,103\,709\,636\,808, \\ s_5 &= 342\,612\,368\,928\,228\,000, \\ s_6 &= 736\,443\,048\,260\,836\,419\,880. \end{aligned}$$

The corresponding complete intersections have the same Pontrjagin numbers and Euler characteristic by [7, §7], so the two complete intersections

$$\begin{aligned} X_6(2323, 2241, 2231, 2117, 2079, 1957, 1953, 1899), \\ X_6(2321, 2263, 2187, 2163, 2037, 2001, 1919, 1909) \end{aligned}$$

are homeomorphic by [4, Theorem 1.1], but have different moduli space dimensions:

$$\begin{aligned} m(\underline{d}) &= 4\,639\,611\,966\,677\,972\,182\,663\,146\,217\,041\,064\,938, \\ m(\underline{d}') &= 4\,639\,610\,187\,986\,885\,926\,979\,324\,513\,081\,980\,800. \end{aligned}$$

Using the method from [8], define the composed multidegree

$$(2.3) \quad d_{\lambda, \mu} = (\underbrace{\underline{d}, \dots, \underline{d}}_{\lambda}, \underbrace{\underline{d}', \dots, \underline{d}'}_{\mu}), \lambda + \mu = s \geq 1.$$

¹The two multidegrees are firstly known as a solution of "Equal Products and Equal Sums of Like Powers", by Chen Shuwen in 2001.

The composed multidegrees $d_{0,s}, d_{1,s-1}, \dots, d_{s,0}$ have the same power sums $s_1, s_2, \dots, s_6$ respectively, so the corresponding complete intersections $X_6(d_{0,s}), X_6(d_{1,s-1}), \dots, X_6(d_{s,0})$ are homeomorphic to each other. We can prove the following inequality:

$$m(d_{\lambda+1,\mu-1}) - m(d_{\lambda,\mu}) > 0, 0 \leq \lambda < s = \lambda + \mu.$$

This inequality will be proved in the coming Proposition.

Now, the sequence $m(d_{\lambda,s-\lambda})|_{\lambda=0,1,\dots,s-1}$ is strictly monotonously increasing. Let $k = s + 1$, there exist k 6-dimensional complete intersections $X_6(d_{0,s}), X_6(d_{1,s-1}), \dots, X_6(d_{s,0})$ in $\mathbb{C}P^{8s+6} = \mathbb{C}P^{8k-2}$ with the desired properties. The proof is finished. $\square$

Proposition 2.2.

$$m(d_{\lambda+1,s-\lambda-1}) - m(d_{\lambda,s-\lambda}) > 0, 0 \leq \lambda < s.$$

Proof. For the chosen multidegrees $\underline{d}$ and $\underline{d}'$ in (2.2),

$$(2.4) \quad m(d_{\lambda,s-\lambda}) = 1 - (N+1)^2 + \left[\lambda \sum_{d_i \in \underline{d}} + (s-\lambda) \sum_{d_i \in \underline{d}'} \right] \binom{N+d_i}{N} \\ - \left[\lambda \sum_{d_i \in \underline{d}} + (s-\lambda) \sum_{d_i \in \underline{d}'} \right] \sum_{\substack{1 \leq k \leq 8s \\ d_k \in d_{\lambda,s-\lambda}}} \binom{N+d_i-d_k}{N},$$

Where, the index j in (2.1) is maximally 1 that is determined by $\max(\underline{d}, \underline{d}') = 2323$ and $\min(\underline{d}, \underline{d}') = 1899$. So,

$$(2.5) \quad m(d_{\lambda+1,s-\lambda-1}) - m(d_{\lambda,s-\lambda}) \\ = \left[\sum_{d_i \in \underline{d}} - \sum_{d_i \in \underline{d}'} \right] \binom{N+d_i}{N} \\ + \left\{ - \left[(\lambda+1) \sum_{d_i \in \underline{d}} + (s-\lambda-1) \sum_{d_i \in \underline{d}'} \right] \sum_{\substack{1 \leq k \leq 8s \\ d_k \in d_{\lambda+1,s-\lambda-1}}} \right. \\ \left. + \left[\lambda \sum_{d_i \in \underline{d}} + (s-\lambda) \sum_{d_i \in \underline{d}'} \right] \sum_{\substack{1 \leq k \leq 8s \\ d_k \in d_{\lambda,s-\lambda}}} \right\} \binom{N+d_i-d_k}{N} \\ := M_0(\lambda, s) + M_1(\lambda, s),$$

Where $M_0(\lambda, s), M_1(\lambda, s)$ are polynomials of invariants s and λ ($N = 8s + 6$):

$$\begin{aligned}
(2.6) \quad M_0(\lambda, s) &= \left[\sum_{d_i \in \underline{d}} - \sum_{d_i \in \underline{d}'} \right] \binom{N + d_i}{N}, \\
M_1(\lambda, s) &= \left[-(\lambda + 1) \sum_{d_i \in \underline{d}} \sum_{d_k \in d_{\lambda+1, s-\lambda-1}} - (s - \lambda - 1) \sum_{d_i \in \underline{d}'} \sum_{d_k \in d_{\lambda+1, s-\lambda-1}} \right. \\
&\quad \left. + \lambda \sum_{d_i \in \underline{d}} \sum_{d_k \in d_{\lambda, s-\lambda}} + (s - \lambda) \sum_{d_i \in \underline{d}'} \sum_{d_k \in d_{\lambda, s-\lambda}} \right] \binom{N + d_i - d_k}{N} \\
(2.7) \quad &= \left[(-2\lambda - 1) \sum_{d_i \in \underline{d}} \sum_{d_k \in \underline{d}} + (1 + 2\lambda - s) \sum_{d_i \in \underline{d}} \sum_{d_k \in \underline{d}'} \right. \\
&\quad \left. + (1 + 2\lambda - s) \sum_{d_i \in \underline{d}'} \sum_{d_k \in \underline{d}} + (2s - 2\lambda - 1) \sum_{d_i \in \underline{d}'} \sum_{d_k \in \underline{d}'} \right] \binom{N + d_i - d_k}{N}.
\end{aligned}$$

From the above, we see that (2.5) is exactly a polynomial of s, λ with complicated coefficients and higher degree. Using the technical computational software **Mathematica**, (2.5) can be computed by executable program (see the code listed in [10, §5]). Finally, we calculate the following results:

$$\begin{aligned}
m(d_{1,0}) - m(d_{0,1}) &= 1\,778\,691\,086\,255\,683\,821\,703\,959\,084\,138, \\
m(d_{2,0}) - m(d_{1,1}) &= 4\,499\,576\,565\,311\,886\,952\,937\,393\,989\,311\,636\,807\,018\,493\,942\,453, \\
m(d_{1,1}) - m(d_{0,2}) &= 4\,499\,576\,565\,312\,040\,117\,972\,354\,794\,044\,912\,596\,557\,706\,541\,183.
\end{aligned}$$

More generally,

$$m(d_{\lambda+1, s-\lambda-1}) - m(d_{\lambda, s-\lambda}) > 10^{65}, 0 \leq \lambda < s, s \geq 3.$$

Thus, it is clear that, with any fixed $s \geq 1, s > \lambda \geq 0$, $m(d_{\lambda, s-\lambda})$ form a strictly monotonously increasing sequence for λ . $\square$

Theorem 2.3. *For each integer $k > 1$, there exist k diffeomorphic complex 7-dimensional complete intersections in $\mathbb{C}P^{15k-8}$ belonging in the moduli space to components with different dimensions.*

Proof. Consider the following two multidegrees([5]²)

$$\begin{aligned}
(2.8) \quad \underline{d} &= (608, 592, 572, 516, 500, 453, 450, 424, 423, 408, 396, 366, 339, 312, 309), \\
\underline{d}' &= (604, 600, 564, 528, 488, 456, 452, 429, 416, 412, 387, 375, 333, 318, 306).
\end{aligned}$$

²These multidegrees are founded by Guo Xianqiang using artificial hand calculation.

They have the same total degree d , powers sums $s_1, \dots, s_7$.

$$\begin{aligned} d &= 3\,753\,247\,176\,539\,885\,786\,786\,848\,165\,802\,803\,200\,000, \\ s_1 &= 6\,668, \\ s_2 &= 3\,094\,964, \\ s_3 &= 1\,495\,641\,932, \\ s_4 &= 749\,415\,139\,508, \\ s_5 &= 387\,496\,273\,524\,068, \\ s_6 &= 205\,753\,667\,680\,942\,844, \\ s_7 &= 111\,680\,899\,229\,310\,068\,732. \end{aligned}$$

Then the corresponding complete intersections

$$\begin{aligned} X_7(608, 592, 572, 516, 500, 453, 450, 424, 423, 408, 396, 366, 339, 312, 309), \\ X_7(604, 600, 564, 528, 488, 456, 452, 429, 416, 412, 387, 375, 333, 318, 306) \end{aligned}$$

are homeomorphic by [4, Theorem 1.1]. Furthermore

$$d = 2^{28} \times 3^{13} \times 5^5 \times 11^2 \times 13^2 \times 17 \times 19 \times 37 \times 43 \times 47 \times 53 \times 61 \times 103 \times 113 \times 151,$$

so they are diffeomorphic by Traving [9] (see also [6, Theorem A]). However, they have different moduli space dimensions:

$$\begin{aligned} m(\underline{d}) &= 44\,406\,795\,197\,386\,326\,965\,368\,167\,342\,722\,355\,968\,367, \\ m(\underline{d}') &= 44\,384\,030\,917\,398\,245\,056\,066\,270\,542\,147\,363\,962\,375. \end{aligned}$$

Similarly, for the composed multidegrees $d_{0,s}, d_{1,s-1}, \dots, d_{s,0}$ (cf. (2.3)), the corresponding complete intersections $X_7(d_{0,s}), X_7(d_{1,s-1}), \dots, X_7(d_{s,0})$ are diffeomorphic to each other.

For the chosen multidegrees $\underline{d}$ and $\underline{d}'$ in (2.8), since $\max(\underline{d}, \underline{d}') = 608$ and $\min(\underline{d}, \underline{d}') = 306$, then

$$\begin{aligned} (2.9) \quad & m(d_{\lambda+1,s-\lambda-1}) - m(d_{\lambda,s-\lambda}) \\ &= \left[\sum_{d_i \in \underline{d}} - \sum_{d_i \in \underline{d}'} \right] \binom{N + d_i}{N} \\ &+ \left\{ - \left[(\lambda + 1) \sum_{d_i \in \underline{d}} + (s - \lambda - 1) \sum_{d_i \in \underline{d}'} \right] \sum_{\substack{1 \leq k \leq 15s \\ d_k \in d_{\lambda+1,s-\lambda-1}}} \right. \\ &\quad \left. + \left[\lambda \sum_{d_i \in \underline{d}} + (s - \lambda) \sum_{d_i \in \underline{d}'} \right] \sum_{\substack{1 \leq k \leq 15s \\ d_k \in d_{\lambda,s-\lambda}}} \right\} \binom{N + d_i - d_k}{N} \\ &:= M_0(\lambda, s) + M_1(\lambda, s), \end{aligned}$$

where $M_0(\lambda, s)$, $M_1(\lambda, s)$ are same as (2.6), (2.7). Finally, we calculate the following results:

$$\begin{aligned}
& m(d_{1,0}) - m(d_{0,1}) \\
&= 22\,764\,279\,988\,081\,909\,301\,896\,800\,574\,992\,005\,992, \\
& m(d_{2,0}) - m(d_{1,1}) \\
&= 33\,455\,700\,664\,468\,562\,980\,578\,980\,033\,713\,637\,615\,501\,603\,407\,170\,478\,458\,745, \\
& m(d_{1,1}) - m(d_{0,2}) \\
&= 33\,455\,700\,663\,954\,152\,769\,609\,839\,164\,207\,754\,699\,356\,499\,185\,723\,378\,335\,507.
\end{aligned}$$

More generally,

$$m(d_{\lambda+1, s-\lambda-1}) - m(d_{\lambda, s-\lambda}) > 3 \times 10^{76}, 0 \leq \lambda < s, s \geq 3.$$

Thus, the sequence $m(d_{\lambda, s-\lambda})|_{\lambda=0,1,\dots,s-1}$ is strictly monotonously increasing. Let $k = s + 1$, there exist k 7-dimensional complete intersections $X_7(d_{0,s}), X_7(d_{1,s-1}), \dots, X_7(d_{s,0})$ in $\mathbb{C}P^{15s+7} = \mathbb{C}P^{15k-8}$ with the desired properties. $\square$

REFERENCES

- [1] Brückmann, P.: A remark on moduli spaces of complete intersections. J. Reine Angew. Math. **476** (1996), 209–215.
- [2] Brückmann, P.: A remark on moduli spaces of 4-dimensional complete intersections. J. Reine Angew. Math. **525** (2000), 213–217.
- [3] Chen, S.W.: Equal products and equal sums of like powers. <http://euler.free.fr/eslp/product.htm>
- [4] Fang, F. Q., Wang, J. B.: Homeomorphism classification of complex projective complete intersections of dimensions 5,6 and 7. Math. Z. **266** (2010), 7919–746.
- [5] Guo, X.Q.: <http://bbs.emath.ac.cn/thread-5853-1-1.html> (in Chinese).
- [6] Kreck, M.: Surgery and duality. Ann. of Math. **149**(3) (1999), 707–754.
- [7] Libgober, A. S., Wood, J. W.: Differentiable structures on complete intersections, I. Topology. **21** (1982), 469–482.
- [8] Libgober, A. S., Wood, J. W.: Remarks on moduli spaces of complete intersections. Contemp. Math. **58** (1986), 183–194.
- [9] Traving, C.: Klassifikation vollständiger Durchschnitte. Diplomarbeit, Mainz (1985). Available at <http://www.mfo.de/Staff/traving.pdf>.
- [10] Wang, J. B.: Remarks on 5-dimensional complete intersections. Electron. Res. Announc. Math. Sci. **21** (2014), 28–40.

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN UNIVERSITY
 WEIJIN ROAD 92, NANKAI DISTRICT, TIANJIN 300072, P.R.CHINA
E-mail address: wjianbo@tju.edu.cn